

Julio Coll

Bull. Eur. Ass. Fish Pathol., 12(5),150, 1992.

NEUTRALISING EPI TOPE(S) OF THE GLYCOPROTEIN OF VIRAL HAEMORRHAGIC SEPTICAEMIA VIRUS ARE EXPRESSED IN THE MEMBRANE OF INFECTED TROUT MACROPHAGES

BY *A. ESTEPA, D.FRÍAS & J.M.COLL

Introduction

To study the *in vitro* VHSV infection of trout macrophages we selected the anterior kidney because it is the major haematopoietic tissue in the rainbow trout, *Oncorhynchus mykiss* (Coll, 1990). Characterisation of the isolated kidney adherent cells as macrophages and melanomacrophages as described previously (Estepa and Coll, 1992a, 1992b). In the fibrin clot *in vitro* system (Coll, 1990; Estepa and Coll, 1992b), cells or mitogen induced colonies from trout kidney were the target of the viral haemorrhagic septicaemia virus (VHSV) (Estepa and Coll, 1991a; Estepa *et al.*, 1991b). It was not clear from those studies whether or not the trout macrophages were also a lytic target for VHSV. Trout macrophages were stimulated by the glycoprotein of the spikes and by the nucleoproteins of the VHSV in cultures of cells from the anterior kidney of healthy trout and of trout immunised by injection with VHSV (Estepa and Coll, 1991c). These results were confirmed by using leucocytes from trout resistant to VHSV one year after infection (Estepa and Coll, 1992a), making clear the importance of the trout macrophages in resistance to VHS. Here we report the *in vitro* conditions under which trout macrophages, presented VHSV neutralising epitopes in their membranes after viral infection.

Material and Methods

Anterior kidney cells from rainbow trout - These cells were prepared as described previously (Coll, 1990; Estepa and Coll, 1992a) from trout of 5-20g body weight. The trout were cooled to 4°C and bled from the tail vein to reduce the blood content of

the kidney. The head kidney (pronephros) was removed and cell suspensions obtained as described. To isolate adherent cells (macrophages), the head kidney cells were incubated in 25cm² bottles during 10 days in the medium described below, then medium was changed and adherent cells kept at 14°C until used for the experiments. The cell culture medium (Flow Labs, Ayrshire, Scotland) was RPMI-1640 (Dutch modification) with 2mM L-glutamine, 1mM sodium pyruvate, 1.2µg/ml amphotericin, 50 µg/ml gentamicin, 20 mM Hepes, 50µM mercaptoethanol, 10% pre-tested foetal calf serum and 0.5% pooled rainbow trout serum.

Virus - The strain of virus used was VHSV 07.71 (gift of Dr.P.de Kinkelin, INRA, Jouy en Josas, France) isolated from rainbow trout. The virus was cultured in epithelioma papillosum cyprine (EPC) cells, and added to the kidney macrophage cultures as supernatant from infected EPC cell monolayers after complete cytopathic effect (Basurco and Coll, 1989).

Immunofluorescence by flow cytometry :- Immunofluorescence of VHSV-infected cells was carried out after macrophages were infected at 0.2 VHSV-07.71 TCID₅₀ per macrophage (cultures containing 1x 10⁶ macrophages/5 ml). Non-infected controls were included in parallel experiments. Incubation was at 14°C for 5 days without 5% CO₂ gassing (Estepa *et al.*, 1991b). Macrophages were detached from the surface of the 25cm² bottles by mechanical agitation and re suspended in PBS (0.05M sodium phosphate, 0.15M sodium chloride, pH7.4) containing 1% bovine serum albumin (BSA) and 0.1% sodium azide. Mechanical agitation separated macrophages from other adherent cells of fibroblastic appearance.





Figure 1.- Effect of VHSV adherence of trout kidney face of 25 cm² bottles (C) then the non-adherent cell again, and infected with anterior kidney cells from size distribution of macro

Macrophage suspensions at 300g for 10 min and pellet was then gently resuspended in BSA, azide containing mouse ascites with the antibodies (Sanz and Coll, 1992a) at 20°C with occasional suspensions were centrifuged in 400-fold anti-mouse IgG-FITC (C. Tilsburg, The Netherlands) for 30 min at 20°C. The suspensions were again centrifuged twice and then resuspended containing 0.3% paraformaldehyde of harvest and staining, 5 were examined by flow

by flow cytometry :- of VHSV-infected but after macrophages VHSV-07.71 TCID₅₀ cultures containing 1x 10⁶. Non-infected controls parallel experiments. Infected for 5 days without 5% (et al, 1991b). Macrophages removed from the surface of by mechanical agitation in PBS (0.05M sodium chloride, pH7.4) and serum albumin (BSA) added. Mechanical agitation of macrophages from other adhesion and plastic appearance.



Becton-Dickinson (San José, California) FACSscan apparatus using the program LYSYS II version 1.0. Green fluorescence was measured at FL1 (514-545 nm).

When the cultures of macrophages were infected with the appropriate titer of VHSV 07.71, total cell lysis occurred after 1 week or more in culture ($n = 6$). Higher multiples of infection resulted in delayed cell lysis or no lysis at all. The macrophage size profile as measured by flow cytometry was as shown in Fig. 1B. After 5 days of VHSV infection the number of larger macrophages decreased with an increase in the number of small macrophages (Fig. 1C). When these

macrophages were stained with neutralising polyclonal or monoclonal (IH10) antibodies, significant differences in fluorescence intensity were found between infected and non-infected macrophages (Fig. 2), whereas

parallel controls with no antibodies present only showed background fluorescence. The infected-cell related immunofluorescence was localised in a macrophage population of small size (Fig. 2).

demonstrated in that showed that the epitope clonal antibodies containing activity and even more neutralising-enhancing activity (Coll, 1992a) are exposed of the infected macrophages. The findings reported here of infected macrophage immune system of the trout are active killers of macrophage because, during development of the disease time to elaborate antibodies also need the macrophages. Further experiments are to investigate these and other

Summary

Macrophages isolated from rainbow trout infected with viral haemorrhagic septicaemia (VHSV) showed positive membrane fluorescence with neutralising polyclonal antibody (MAb) anti G IH10. The possibility of using trout macrophages to study the epitopes related against VHSV.

Acknowledgements

We acknowledge the technical assistance of J. Coll Perez in typing. This work was supported by Research Grant (8568) from the Ministerio de Investigaciones Agrarias de España (Spain).

Discussion

The importance of macrophages (monocytes) as accessory cells in higher vertebrate immune responses is well established, but their function in fish is not yet completely understood (Estepa *et al*, 1991c; Estepa and Coll, 1992a). Because of the possibility of obtaining trout macrophage populations

with a high degree of purification by plastic adherence, these could easily be infected *in vitro* with VHSV and their VHSV epitope expression studied by flow cytometry. Although the average size of the macrophage population decreased after infection, before complete lysis occurred, positive membrane fluorescence could be






Figure 2. Flow cytometry of macrophages stained with MAb IH10 and with PAb. —A—, VHSV infected; —B—, non-infected. IH10, MAb anti-glycoprotein of VHSV with neutralising-enhancing activity (Sanz and Coll, 1992). PAb, polyclonal antibody obtained from the mouse used to make the anti-VHSV hybridomas.

no antibodies present
und fluorescence. The
immunofluorescence
macrophage population

PAb



with PAb.
protein of VHSV with
monoclonal antibody obtained

f purification by plastic
ld easily be infected in
d their VHSV epitope
by flow cytometry.
verage size of the
tion decreased after in-
complete lysis occurred,
fluorescence could be

demonstrated in that population. This showed that the epitopes defined by polyclonal antibodies containing neutralising activity and even more by MAb IH10 with neutralising-enhancing activity (Sanz and Coll, 1992a) are exposed in the membrane of the infected macrophages.

The findings reported here open the possibility of further studies into the recognition of infected macrophages by cells of the immune system of the trout. Most probably, some of the cells from VHSV resistant trout are active killers of any infected trout macrophage because, due to the fast development of the disease there is almost no time to elaborate antibodies which would also need the macrophages to be elaborated. Further experiments are in progress to investigate these and other possibilities.

Summary

Macrophages isolated from rainbow trout kidney and infected with viral haemorrhagic septicaemia virus (VHSV) showed positive membrane immunofluorescence with neutralising polyclonal and monoclonal antibody (MAb) anti G IH10. These findings open the possibility of using trout macrophages as presenting cells to study the epitopes relevant to the protection against VHSV.

Acknowledgements

We acknowledge the technical assistance of P. Parrilla and of J. Coll Perez in typing. This work was supported by Research Grant (8568) from the Instituto Nacional de Investigaciones Agrarias del Ministerio de Agricultura (Spain).

Authors Address

INIA-Sanidad Animal. Embajadores, 68 28012 - Madrid, Spain.

*To whom correspondence should be sent.

References

- Basurco, B. and Coll, J.M. (1989). Spanish isolates and reference strains of viral haemorrhagic septicaemia virus show similar protein size patterns. Bull. Eur. Ass. Fish. Pathol., 9, 92-95.
- Coll, J.M. (1990). Estimulación de colonias de células de riñón de trucha con fitohemaglutinina en cultivos de fibrina. Inmunología, 9, 140-145.
- Estepa, A. and Coll, J. M. (1991a). Infection of mitogen stimulated colonies from trout kidney cell cultures with salmonid viruses. J. Fish Dis. 14, 555-562.
- Estepa, A., Frías, D. and Coll, J.M. (1991b). Infection of trout kidney cells with infectious pancreatic necrosis and viral haemorrhagic septicaemia viruses. Bull Eur. Ass. Fish Dis. 11, 101-104.
- Estepa, A., Basurco, B., Sanz, F. and Coll, J.M. (1991c). Stimulation of adherent cells by the addition of purified proteins of viral haemorrhagic septicaemia virus to trout kidney cell cultures. Viral Immunol., 4, 43-52.
- Estepa, A. and Coll, J.M. (1992a). In vitro immunostimulants for optimal responses of kidney leucocytes from trout surviving viral haemorrhagic septicaemia virus disease. J. Fish and Shellfish Immunol., 2, 53-68.
- Estepa, A. and Coll, J.M. (1992b). Mitogen induced proliferation of trout kidney leucocytes in fibrin clots. Vet. Immunol. Immunopathol. (in press).
- Sanz, F., and Coll, J.M. (1992a). Neutralising enhancing monoclonal antibody recognises the denatured glycoprotein of the viral haemorrhagic septicaemia virus (VHSV). Archiv. Virol. (in press).
- Sanz, F., Basurco, B., Babin, M., Domínguez, J. and Coll, J.M. (1992b). Viral haemorrhagic septicaemia virus isolates studied with monoclonal antibodies against its structural proteins. J. Fish Dis. (submitted).